

PREDATION ON THE COASTAL TAILED FROG (*ASCAPHUS TRUEI*)
BY A SHREW (*SOSEX SPP.*) IN WASHINGTON STATE

ERIC M LUND, MARC P HAYES, TIERRA R CURRY, JEFFREY S MARSTEN, AND
KEVIN R YOUNG

Key words: Tailed Frog, *Ascaphus truei*, Water Shrew, *Sorex palustris*, Marsh Shrew, *Sorex bendirii*, predation, Washington State

Few observations exist of predation on post-metamorphic Coastal Tailed Frogs (*Ascaphus truei*). Karraker (2001) found a Common Garter Snake (*Thamnophis sirtalis*) with an adult male *A. truei* (36 mm snout-vent length [SVL]) in its mouth, and predation of metamorphosing *A. truei* by a hellgrammite (Megaloptera) and a larval Cope's Giant Salamander (*Dicamptodon copei*) also have been reported (Jones and Raphael 1998). To our knowledge, no reports exist of predation on *A. truei* by mammals. Here we report an observation of *A. truei* being captured and eaten by a shrew (*Sorex spp.*).

At 12:15 on 17 May 2007, JSM captured an adult female *A. truei* (approximately 40 mm SVL) 2 m from a stream edge along an unnamed tributary of the West Fork Rue Creek, Pacific County, Washington (5160283N, 445091E, UTM Zone 10, Datum WGS 1984; elevation 282 m). The capture site is along a 2nd-order stream segment (based on Strahler 1952) with a mean wetted channel width of 1.3 m at

the time of this observation; and located 220 m upstream of the electroshock-determined end of fish-bearing waters (Jason Walters, Weyerhaeuser Company, Federal Way, WA, pers. comm.) and 150 m below the stream origin. The site was located in a 60-y-old stand of Western Hemlock (*Tsuga heterophylla*) with a closed canopy. Understory vegetation included a scattered shrub layer of Oval-leaved Huckleberry (*Vaccinium ovalifolium*), Fool's Huckleberry (*Menziesia ferruginea*), and ground cover of mostly Wood Sorrel (*Oxalis oregana*), Deer Fern (*Blechnum spicant*), and Sword Fern (*Polystichum munitum*). The female *A. truei* was handled for measurement for about 2 min prior to being returned to a small stream pool (0.5 × 0.7 m, maximum depth 15 cm).

About 5 min later, EML, JSM, TRC, and KRY observed a large, black shrew with a whitish underside moving in and out of woody debris along the stream bank. The shrew came into full view, swam across the stream, passing within 0.5 m of the frog that had remained suspended at pool surface near its original release site, and then paused on the opposite stream

bank approximately 1 m away from the frog. It then pivoted, re-entered the stream and lunged at the *A. truei*, capturing it headfirst with its forelimbs and mouth. It swam to the edge of the stream with the frog, which remained peculiarly motionless in its mouth. It then grasped the *A. truei* between its front limbs and began chewing on it without taking cover. After about 10 s of chewing the shrew moved out of sight into a small hole in the stream bank with the frog in its mouth.

Without having the shrew in hand, we were not able to confirm the species identity. The Northern Water Shrew (*Sorex palustris*) and Marsh Shrew (*S. bendirii*), the 2 largest species of *Sorex* in North America, are both described as having black dorsal pelage and are often found close to small, forested streams in Washington State (Pattie 1973; Beneski and Stinson 1987; Eder 2002). Though the location of this observation might suggest *S. bendirii* (Ingles 1965; Pattie 1973), the whitish underside is more typical of *S. palustris* (Beneski and Stinson 1987; Eder 2002).

Shrew predation on other stream-associated amphibian species has been documented. Predation on *Dicamptodon* spp. by *S. palustris* has been reported several times (for example, Sorenson 1962; Nussbaum and Maser 1969; Beneski and Stinson 1987), and salamander remains have been found in the stomachs of various shrew species (Hamilton 1930); but shrew predation on anurans in the wild has only rarely been observed. Maier (2005) made 2 field observations of presumed predatory behavior by the Northern Short-Tailed Shrew (*Blarina brevicauda*) towards the Eastern Spadefoot (*Scaphiopus holbrookii*), but in both cases the observer disturbed the shrew, and no injuries were found on the toads. To our knowledge, the only report of shrew predation on an anuran in the wild is that of Dharmakumarsinhji (1946) who reported an attack on an Indian Bullfrog (*Rana tigrina*) by a Musk-shrew (*Suncus caeruleus*). Although reports of shrews preying on tailed frogs (*Ascaphus* spp.) are lacking, the habitats along forested streams typically used by *S. palustris* and *S. bendirii* (Pattie 1973, Beneski and Stinson 1987; Anthony and others 1987) overlap substantially with habitats used by *A. truei* (Diller and Wallace 2001; Nielsen and others 2001). Shrews may prey on tailed frogs in the

Pacific Northwest more frequently than our isolated observation suggests.

Acknowledgments.—Fieldwork was supported by the Forests and Fish Adaptive Management Program, Washington Department of Fish and Wildlife (WDFW). Washington Department of Natural Resources facilitated work on their land, where this observation was made. Studies were done under a programmatic WDFW handling permit provided to employees in the course of their work; conditions of this permit require adherence to the guidelines for use of live amphibians and reptiles in the field (Beaupre and others 2004). This is contribution No. 22 of the Forests and Fish Section of the WDFW Habitat Program Amphibian Research Group.

LITERATURE CITED

- ANTHONY RG, FORSMAN ED, GREEN GA, WITMER G, NELSON SK. 1987. Small mammal populations in riparian zones of different-aged coniferous Forests. *The Murrelet* 68:94–102.
- BEAUPRE SB, JACOBSON ER, LILLYWHITE HB, ZAMUDIO K. 2004. Guidelines for use of live amphibians and reptiles in field and laboratory research, second edition. Herpetological Animal Care and Use Committee of the American Society of Ichthyologists and Herpetologists.
- BENESKI JR JT, STINSON DW. 1987. *Sorex palustris*. *Mammalian Species* 296:1–6.
- DHARMAKUMARSINHJI KS. 1946. Musk shrew (*Suncus caeruleus*) attacking bull-frog (*Rana tigrina*). *The Journal of the Bombay Natural History Society* 46: 180–181.
- DILLER LV, WALLACE RL. 1999. Distribution and habitat of *Ascaphus truei* in streams in managed, young growth forests in North Coastal California. *Journal of Herpetology* 33:71–79.
- EDER T. 2002. *Mammals of Washington and Oregon*. Renton, WA: Lone Pine Publishing. 351p.
- HAMILTON JR WJ. 1930. The food of Soricidae. *Journal of Mammalogy* 11:26–39.
- INGLES L. 1965. *Mammals of the Pacific States*. Stanford, CA: Stanford University Press. 506p.
- JONES LC, RAPHAEL M. 1998. *Ascaphus truei* (Tailed Frog). *Predation*. *Herpetological Review* 29:39.
- KARRAKER N. 2001. *Ascaphus truei* (Tailed Frog). *Predation*. *Herpetological Review* 32:100.
- MAIER T. 2005. Predatory behaviors of *Blarina brevicauda* toward a fossorial Eastern Spadefoot toad (*Scaphiopus holbrookii*). *Special Publication of the International Society of Shrew Biologists* 1:361–366.
- NIELSEN M, LOHMAN K, SULLIVAN J. 2001. Phylogeography of the Tailed Frog (*Ascaphus truei*): Implications for the biogeography of the Pacific Northwest. *Evolution* 55:147–160.
- NUSSBAUM RA, MASER C. 1969. Observations of *Sorex*

- palustris* preying on *Dicamptodon ensatus*. The Murrelet 50:23–24.
- PATTIE D. 1973. *Sorex bendirii*. Mammalian Species 27:1-2.
- SORENSEN MW. 1962. Some aspects of water shrew behavior. American Midland Naturalist 68:445–462.
- STRAHLER AN. 1952. Hypsometric (area-altitude) analysis of erosional topography. Bulletin Geological Society of America 63:1117–1142.
- Washington Department of Fish and Wildlife, Habitat Program, 600 Capital Way North, Olympia, Washington 98501, USA; lundeml@dfw.wa.gov (EML, MPH); Center for Biological Diversity, PO Box 11374, Portland, Oregon 97202 (TRC); 617 SW 135th St, Burien, WA 98146 (JSM); 1617 Quince St NE, Olympia, WA 98506 (KRY). Submitted 09 May 2008, accepted 23 June 2008. Corresponding Editor: Paul Cryan.